

THE MILDLY METAL-POOR GLOBULAR CLUSTER M4: STAR-TO-STAR ABUNDANCE VARIATIONS IN 20 GIANTS

I. I. IVANS AND C. SNEDEN

McDonald Observatory, University of Texas, Austin, USA

AND

R.P. KRAFT

Lick Observatory, University of California, Santa Cruz, USA

Many low-metallicity globular clusters exhibit large star-to-star variations of C, N, O, Na, Mg, & Al abundances. In higher-metallicity clusters, the abundance swings are muted. Evidence of anticorrelations of O versus Na and Mg versus Al exists in classical northern-hemisphere clusters spanning a range of metallicities, $-0.8 \leq [\text{Fe}/\text{H}] \leq -2.2$. These and other anticorrelations, observed to be a function of giant branch position, are consistent with material having undergone proton-capture nucleosynthesis, presumably taking place in regions where the CNO-cycle is in operation, and brought to the surface by a deep-mixing mechanism. Distinctly bimodal distributions of cyanogen strengths at nearly all giant branch positions are observed in some clusters, including M4 (Norris 1981). The metallicity of M4 ($[\text{Fe}/\text{H}] \sim -1$) places it among clusters in which the O versus Na and Mg versus Al anticorrelations are expected to be suppressed. Indeed, not much abundance variation has been observed in small samples of M4 giant stars (see e.g. Brown & Wallerstein 1992). This puzzle led us to consider an abundance study of a large sample of bright giants in the mildly metal-poor globular cluster M4, the nearest, brightest, and one of the most accessible targets in which to study the CN-bimodal phenomenon.

We observed 20 M4 giant stars in 1997 using high resolution échelle spectrographs covering the effective wavelength range: $\lambda\lambda$ 5200 to 8800 Å on the 2.7-m at McDonald Observatory ($R \approx 60,000$) and the 3.0-m at Lick Observatory ($R \approx 50,000$).

Large and differential interstellar reddening due to the dark nebulosity in Scorpio-Ophiuchus does not permit a careful estimate of reddenings for individual stars; one cannot assume a strict $(B-V)$ versus T_{eff} relationship. Instead, we obtained precise T_{eff} s by employing two different spectroscopic

techniques: (1) ratios of central depths of temperature sensitive absorption features (Gray 1994) to initially estimate relative stellar Teffs and (2) as in previous Lick-Texas work, final Teffs were determined in conjunction with surface gravity and microturbulence constraints.




Figure 1. Boxplot of M4 Giant Star Abundances - A boxed horizontal line indicates the interquartile range [IQ] & median found for a particular element. Vertical tails extending from the boxes indicate the total range of abundances determined for each element, excluding outliers. Mild outliers (within $1.5 \times \text{IQ}$) are denoted by hollow circles (o) & severe outliers (from 1.5 to $3 \times \text{IQ}$) by filled circles ($\bullet$). The dashed line represents the solar value for a particular elemental abundance.

The scatter about the mean (of elements not expected to be sensitive to proton-capture nucleosynthesis) compares well to those obtained in other high resolution cluster work and, as expected, no trend with Teff is observed for these non-volatiles. In individual M4 stars, anticorrelated abundances of N/Na/Al with C/O provide strong evidence for high-temperature proton-capture nucleosynthesis. We also observe increasing O-depletion/N-enhancement as stars ascend the RGB but combined $\log_{\epsilon}(\text{C+N+O})$ remains $\sim$ constant. Extremely low carbon isotope ratios ($^{12}\text{C}/^{13}\text{C} = 4.6 \pm 0.8$) indicate that the onset of mixing of the CNO-cycled material occurs at a luminosity lower than that of our observed stars. Elements Na, Mg, Si, & Ca are all enhanced with respect to the iron abundance, in approximate agreement with previous high resolution studies of M4 (Brown & Wallerstein, 1992). Al and Ba show uniform enhancements indicative of primordial enrichment. The analysis also confirms anomalous absorption properties of dust along line of sight to M4 ($E(V - K)/E(B - V) = R = 3.8 \pm 0.4$) that deviate from the normal law of interstellar extinction.

References

- Brown, J. A. & Wallerstein, G. (1992) *Astronomical Journal*, **104**, pp. 1818–1829.
 Gray, D. F. (1994) *PASP*, **106**, pp. 1248–1257.
 Norris, J. E. (1981) *Astrophysical Journal*, **248**, pp. 177–188.